

Exact S matrices and extended supersymmetry

E. Abdalla^{*}, M.C.B. Abdalla[†]

CERN-Theory Division
1211 Geneve 23, Switzerland.

Abstract

We study the constraint imposed by supersymmetry on the exact S -matrix of $\mathbb{C}P^{n-1}$ model, and compute a non-trivial phase factor in the relation between the S -matrix and one of the supersymmetry generators. We discuss several features connected with the physical interpretation of the result. The supersymmetry current is studied in such context as well, and we find some operators appearing in the conservation equation of the supersymmetry current. The relation with the literature on the subject is also discussed.

^{*} Work supported by FAPESP. Permanent address: Instituto de Física - USP, C.P. 20516, S.Paulo, Brazil.

[†] Supported by CNPq. Permanent address: Instituto de Física Teórica - UNESP, R. Pamplona 145, CEP-01405, S.Paulo, Brazil.

The issue of supersymmetry has been playing a central role in the development of quantum field theory since two decades. Supersymmetric grand unified theories are the most promising candidates to extend the standard model of strong and electro-weak interactions. In the string domain supersymmetry is essential in order to define a tachyon free theory. Nevertheless, it must be broken already at intermediate energies, in order to take into account the observed particle spectrum as revealed from experiments.

Integrable theories on the other hand offer an extraordinary theoretical laboratory to test ideas in quantum field theory. They provide examples of models which are exactly soluble at the level of S -matrix. Thus the analysis of supersymmetric integrable theories is worthwhile to pursue in order to analyse ideas.

On the other hand, there has been recently a large number of results concerning integrable $N = 2$ supersymmetric models. Gepner's conjecture¹, linking these models to perturbations of superconformally invariant theories with Landau-Ginzburg potentials have been put forward by a number of authors^{2,3}. For the time being, this program has been given quite a solid basis, relying on results from singularity (catastrophe) theory^{4,5} on the one hand, in order to obtain a classification of the Landau-Ginzburg theories, and computations based on the thermodynamic Bethe ansatz obtaining⁶ the central charge associated with the exact S -matrices of the given $N = 2$ supersymmetric theory, on the other hand.

The afore mentioned relations lead to a complete classification of the integrable $N = 2$ supersymmetric theories in terms of perturbed Landau-Ginzburg potentials⁷.

In this spirit, a number of examples have been analysed. The supersymmetric generalizations of coset space models - or Kazama Suzuki models⁸, have been studied. These coset models are characterized in terms of a cohomology for the elements of the chiral primary rings⁷, and some models are indeed represented as Landau-Ginzburg theories. Fendley and Intriligator⁶ considered minimal theories perturbed by the least relevant operator. They considered the peculiar phenomenon of fractional fermion number in two-dimensions, where in polymer systems a fermion can be distorted into n solitons, which carry, as a consequence, fermion number $1/n$. This sort of phenomenon plays a central role in our discussion. In fact, they stated that the fractional fermion number is crucial for obtaining the correct soliton content and S -matrix.

The S -matrix corresponding to a $Z(n)$ supersymmetric theory has also been computed in [3]. A previous calculation of the same S -matrix was given by [9], and both results are different. The latter did not contain supersymmetry as a constraint for the S -matrix.

We do not discuss whether any of these results is either correct or incorrect, since from our point of view, a dynamical principle is required to choose which of them will describe the given physics. Indeed, a $Z(n)$ invariant model displaying the S -matrix given by [9] has to display a breakdown of supersymmetry, at least if it is to be realized locally. On the other hand, the results from [3] are supersymmetric, and must describe models which are supersymmetric at the quantum level.

Here we consider a similar problem for the computation of the exact S -matrix of the supersymmetric $\mathbb{C}P^{n-1}$ model. In this case, we verify that the S -matrix fulfills the supersymmetry constraint, only if one considers carefully certain phase factors associated with the statistics of the asymptotic fields. We first make several consistency checks of the

proposed S -matrix, as well as the $1/n$ perturbative check. We consider in detail the case of $\mathbb{CP}^1$ and compare it with the $O(3)$ S -matrix, and we see that the equivalence of $\mathbb{CP}^1$ and $O(3)$ models, verified in the purely bosonic case at the S -matrix level¹⁰, as well as by means of numerical simulations¹¹ seems to break down in the supersymmetric case. We shall discuss the consequences and possible causes of such behavior, as well as whether one can argue in favor of a supersymmetry anomaly in the $\mathbb{CP}^{n-1}$ models.

We have computed some candidates to a supersymmetry anomaly in the context of the $1/n$ expansion of the model. Whether these candidates are anomalies or not can only be decided by the study of the cohomology of such operators¹². Further explanations of the results obtained may be a failure in the description of asymptotic states, or a non standard action of the supersymmetry generators.

The supersymmetric $\mathbb{CP}^{n-1}$ model is defined by the Lagrangian density¹³

$$\mathcal{L} = \overline{D_\mu z} D^\mu z + \overline{\psi} i \not{D} \psi + \frac{f}{2n} (\overline{\psi} \gamma_\mu \psi)^2 + \frac{f}{2n} \left[(\overline{\psi} \psi)^2 - (\overline{\psi} \gamma_5 \psi)^2 \right] , \quad (1)$$

with the constraints $\overline{z}_i z_i = \frac{n}{2f}$ and $\overline{z}_i \psi_i = 0 = \overline{\psi}_i z_i$, where $D_\mu = \partial_\mu - \frac{2f}{n} \overline{z} \partial_\mu z$.

This Lagrangian is invariant under the supersymmetry transformation

$$\begin{aligned} \delta z_i &= \overline{\epsilon} \psi_i , \\ \delta \psi_i &= -i \not{D} z_i \epsilon + \frac{1}{2} z_i \left[\overline{\psi} \psi + \overline{\psi} \gamma_5 \psi \gamma_5 + \overline{\psi} \gamma^\mu \psi \gamma_\mu \right] \epsilon , \end{aligned} \quad (2)$$

where ϵ is a Grassmann valued parameter. This symmetry will be the central issue of the present discussion. The $1/n$ expansion of this model is well known and will not be discussed here. Further details will be given in a more detailed publication.

We consider the exact S -matrix of the supersymmetric $\mathbb{CP}^{n-1}$ model. We make the ansatz (the backward particle-antiparticle scattering vanishes as a consequence of the conservation of the non-local charges; we have already used this fact in the ansatz¹⁴)

$$\begin{aligned} \langle z_\beta(\theta'_1) z_\delta(\theta'_2) | z_\alpha(\theta_1) z_\gamma(\theta_2) \rangle &= [v_1(\theta) \delta_{\alpha\beta} \delta_{\gamma\delta} + v_2(\theta) \delta_{\alpha\delta} \delta_{\gamma\beta}] \delta(\theta_1 - \theta'_1) \delta(\theta_2 - \theta'_2) \\ &\quad + [v_1(\theta) \delta_{\alpha\delta} \delta_{\gamma\beta} + v_2(\theta) \delta_{\alpha\beta} \delta_{\gamma\delta}] \delta(\theta_1 - \theta'_2) \delta(\theta_2 - \theta'_1) , \\ \langle \psi_\beta(\theta'_1) \psi_\delta(\theta'_2) | \psi_\alpha(\theta_1) \psi_\gamma(\theta_2) \rangle &= [u_1(\theta) \delta_{\alpha\beta} \delta_{\gamma\delta} + u_2(\theta) \delta_{\alpha\delta} \delta_{\gamma\beta}] \delta(\theta_1 - \theta'_1) \delta(\theta_2 - \theta'_2) \\ &\quad - [u_1(\theta) \delta_{\alpha\delta} \delta_{\gamma\beta} + u_2(\theta) \delta_{\alpha\beta} \delta_{\gamma\delta}] \delta(\theta_1 - \theta'_2) \delta(\theta_2 - \theta'_1) , \\ \langle \psi_\beta(\theta'_1) z_\delta(\theta'_2) | \psi_\alpha(\theta_1) z_\gamma(\theta_2) \rangle &= [c_1(\theta) \delta_{\alpha\beta} \delta_{\gamma\delta} + c_2(\theta) \delta_{\alpha\delta} \delta_{\gamma\beta}] \delta(\theta_1 - \theta'_1) \delta(\theta_2 - \theta'_2) \\ &\quad + [d_1(\theta) \delta_{\alpha\delta} \delta_{\gamma\beta} + d_2(\theta) \delta_{\alpha\beta} \delta_{\gamma\delta}] \delta(\theta_1 - \theta'_2) \delta(\theta_2 - \theta'_1) , \quad (3) \\ \langle z_\beta(\theta'_1) \overline{z}_\delta(\theta'_2) | z_\alpha(\theta_1) \overline{z}_\gamma(\theta_2) \rangle &= [v_1(i\pi - \theta) \delta_{\alpha\beta} \delta_{\gamma\delta} + v_2(i\pi - \theta) \delta_{\alpha\gamma} \delta_{\delta\beta}] \delta(\theta_1 - \theta'_1) \delta(\theta_2 - \theta'_2) , \\ \langle \psi_\beta(\theta'_1) \overline{\psi}_\delta(\theta'_2) | \psi_\alpha(\theta_1) \overline{\psi}_\gamma(\theta_2) \rangle &= [u_1(i\pi - \theta) \delta_{\alpha\beta} \delta_{\gamma\delta} + u_2(i\pi - \theta) \delta_{\alpha\gamma} \delta_{\delta\beta}] \delta(\theta_1 - \theta'_1) \delta(\theta_2 - \theta'_2) , \\ \langle z_\beta(\theta'_1) \overline{z}_\delta(\theta'_2) | \psi_\alpha(\theta_1) \overline{\psi}_\gamma(\theta_2) \rangle &= [d_1(i\pi - \theta) \delta_{\alpha\beta} \delta_{\gamma\delta} + d_2(i\pi - \theta) \delta_{\alpha\gamma} \delta_{\delta\beta}] \delta(\theta_1 - \theta'_1) \delta(\theta_2 - \theta'_2) , \\ \langle z_\beta(\theta'_1) \overline{\psi}_\delta(\theta'_2) | z_\alpha(\theta_1) \overline{\psi}_\gamma(\theta_2) \rangle &= [c_1(i\pi - \theta) \delta_{\alpha\beta} \delta_{\gamma\delta} + c_2(i\pi - \theta) \delta_{\alpha\gamma} \delta_{\delta\beta}] \delta(\theta_1 - \theta'_1) \delta(\theta_2 - \theta'_2) . \end{aligned}$$

Proceeding as usually, one defines the action of the quantum non-local charge on asymptotic states, and one arrives at the solution

$$v_1(\theta) = \frac{\sin\left(\frac{\theta}{2i} - \frac{\pi}{n}\right)}{\sin \frac{\theta}{2i}} c_1(\theta) , \quad u_1(\theta) = \frac{\sin\left(\frac{\theta}{2i} + \frac{\pi}{n}\right)}{\sin \frac{\theta}{2i}} c_1(\theta) , \quad d_1(\theta) = -\frac{\sin\left(\frac{\pi}{n}\right)}{\sin \frac{\theta}{2i}} c_1(\theta) , \quad (4)$$

and

$$\begin{aligned} v_2(\theta) &= \frac{2i\pi}{n\theta} v_1(\theta) \quad , \quad u_2(\theta) = \frac{2i\pi}{n\theta} u_1(\theta) \quad , \\ c_2(\theta) &= -\frac{2i\pi}{n\theta} c_1(\theta) \quad , \quad d_2(\theta) = -\frac{2i\pi}{n\theta} d_1(\theta) \quad , \end{aligned} \quad (5)$$

which contains also the bound state structure of the model as we will comment later. Equations (5) are a consequence of the non-local conservation laws, while (4) are a consequence of factorization of the amplitude for the scattering of the two bosons and one fermion, as well as crossing. Finally, we determine $c_1(\theta)$ from unitarity.

$$c_1(\theta) = \prod_{l=0}^{\infty} \frac{\Gamma\left(\frac{\theta}{2\pi i} + \frac{1}{n} + l\right) \Gamma\left(1 - \frac{\theta}{2\pi i} + l\right) \Gamma\left(\frac{\theta}{2\pi i} - \frac{1}{n} + l\right) \Gamma\left(1 - \frac{\theta}{2\pi i} + l\right)}{\Gamma\left(1 - \frac{\theta}{2\pi i} + \frac{1}{n} + l\right) \Gamma\left(\frac{\theta}{2\pi i} + l\right) \Gamma\left(1 - \frac{\theta}{2\pi i} - \frac{1}{n} + l\right) \Gamma\left(\frac{\theta}{2\pi i} + l\right)} \quad . \quad (6)$$

This shows the existence of bound states, with the spectrum given by the fusion rule by

$$m_l = m \frac{\sin \frac{l\pi}{n}}{\sin \frac{\pi}{n}} \quad , \quad l = 1, \dots, n-1 \quad . \quad (7)$$

In this theory, a bound state of $n-1$ fermions is equivalent to a antiboson, i.e.

$$\bar{z}_\delta = \frac{1}{(n-1)!} \epsilon_{\delta\alpha_1 \dots \alpha_{n-1}} \psi_{\alpha_1} \dots \psi_{\alpha_{n-1}} \quad , \quad (8)$$

while the bound state of $n-2$ fermions and a boson is a antifermion

$$\bar{\psi}_\alpha = \frac{1}{(n-1)!} \epsilon_{\alpha\beta_1 \dots \beta_{n-1}} \psi_{\beta_1} \dots \psi_{\beta_{n-2}} z_{\beta_{n-1}} \quad . \quad (9)$$

From these formulae it follows that the particle-antiparticle backward scattering amplitudes vanish identically. Using the $1/n$ expansion we can check the leading terms in the above amplitudes (see [14] for details).

Let us consider the ansatz (3) for the S -matrix. The action of the supersymmetry charge on asymptotic states is easy to obtain. We shall only consider the action of the supersymmetry charge in one sector, in such a way that complications arising from topological sectors¹⁵ are immediately dismissed. See also [3] and [6]. We shall come back to this point later. The action of the charge on a two particle state is given by

$$\begin{aligned} Q|z_i(\theta_1)z_j(\theta_2)\rangle &= e^{\frac{\theta}{4}}|\psi_i(\theta_1)z_j(\theta_2)\rangle + e^{i\epsilon}e^{-\frac{\theta}{4}}|z_i(\theta_1)\psi_j(\theta_2)\rangle \quad , \\ Q|\psi_i(\theta_1)z_j(\theta_2)\rangle &= e^{\frac{\theta}{4}}|z_i(\theta_1)z_j(\theta_2)\rangle + e^{i\epsilon'}e^{-\frac{\theta}{4}}|\psi_i(\theta_1)\psi_j(\theta_2)\rangle \quad , \end{aligned} \quad (10)$$

where ϵ and ϵ' are related with the fermionic content of the particles z_i and ψ_j . Usually one expects $\epsilon = 0$ and $\epsilon' = \pi$.

We need also the S -matrix elements

$$\begin{aligned} S|z_i(\theta_1)z_j(\theta_2)\rangle &= (v_1(\theta)\delta_{ik}\delta_{jl} + v_2(\theta)\delta_{il}\delta_{jk})|z_l(\theta_2)z_k(\theta_1)\rangle \quad , \\ S|\psi_i(\theta_1)z_j(\theta_2)\rangle &= (c_1(\theta)\delta_{ik}\delta_{jl} + c_2(\theta)\delta_{il}\delta_{jk})|z_l(\theta_2)\psi_k(\theta_1)\rangle + \\ &\quad + (d_1(\theta)\delta_{il}\delta_{jk} + d_2(\theta)\delta_{ik}\delta_{jl})|\psi_k(\theta_2)z_l(\theta_1)\rangle \quad . \end{aligned} \quad (11)$$

We analyse the commutator of the S -matrix with the supersymmetry charge Q , that is, we impose the condition

$$\langle z_k(\theta'_1)\psi_l(\theta'_2)|QS|z_i(\theta_1)z_j(\theta_2)\rangle = \langle z_k(\theta'_1)\psi_l(\theta'_2)|SQ|z_i(\theta_1)z_j(\theta_2)\rangle \quad . \quad (12)$$

The above equation leads to very simple constraints on the S -matrix elements (11). We are led to the equations

$$\begin{aligned} e^{i\epsilon}e^{\frac{\theta}{4}}v_1(\theta) &= e^{\frac{\theta}{4}}c_1(\theta) + e^{i\epsilon}e^{-\frac{\theta}{4}}d_1(\theta) \quad , \\ e^{\frac{\theta}{4}}v_2(\theta) &= e^{\frac{\theta}{4}}c_2(\theta) + e^{i\epsilon}e^{-\frac{\theta}{4}}d_1(\theta) \quad , \\ e^{-\frac{\theta}{4}}v_1(\theta) &= e^{\frac{\theta}{4}}d_1(\theta) + e^{i\epsilon}e^{-\frac{\theta}{4}}c_1(\theta) \quad , \\ e^{-\frac{\theta}{4}}v_2(\theta) &= e^{\frac{\theta}{4}}d_2(\theta) + e^{i\epsilon}e^{-\frac{\theta}{4}}c_2(\theta) \quad . \end{aligned} \quad (13)$$

The question which arises is whether such equations are compatible with the solution given in eqs. (3-5). If we consider the relation

$$e^{i\epsilon}e^{\frac{\theta}{2}}\frac{e^{i\epsilon}v_1(\theta) - c_1(\theta)}{d_1(\theta)} = -e^{i\epsilon}e^{\frac{\theta}{2}}\frac{e^{i\epsilon}\sin\left(\frac{\theta}{2i} - \frac{\pi}{n}\right) - \sin\frac{\theta}{2i}}{\sin\frac{\pi}{n}} \quad , \quad (14)$$

we immediately see that it is not one as required by the first equation in (14) unless $\epsilon = \frac{i\pi}{n}$! On the other hand, we will see that the candidate to a supersymmetry anomaly is proportional to $\phi_5\gamma_5c$. Witten¹³ has already proven long ago, that ϕ_5 differs by multiples of $2\pi/n$ between “in” and “out” states. Therefore, the anomaly found in order $1/n$ can be interpreted as a result of the above equation relating the “in” and “out” states.

Also connected with this question is the fact that, as shown in [14], the $N = 2$ supersymmetry algebra contains central terms due to the soliton content of the theory. Notice however, that we have only used the $N = 1$ subalgebra. In such a case, we can extract, out of the full algebra

$$\{Q'_\alpha, Q'_\beta\} = \{\overline{Q}'_\alpha, \overline{Q}'_\beta\} = 0 \quad , \quad (15)$$

$$\{Q'_\alpha, \overline{Q}'_\beta\} = \gamma^\mu_{\alpha\beta}P_\mu + \delta_{\alpha\beta}\phi + \gamma_{5\alpha\beta}\phi_5 \quad , \quad (16)$$

the subalgebra

$$Q_+ = Q'_2 + Q'^\dagger_2 = Q'_2 + (\overline{Q}\gamma^0)_2 \quad , \quad (17)$$

such that $Q_+^2 = P_+$; indeed, we have only considered Q_+ . The question about the $U(1)$ charge carried by the supersymmetry charge has to be addressed at this point. In fact,

from the physical interpretation of the model, and especially from the bound state structure (7-9), one learns that the elementary fields carry a fractional extra charge equal to $\pm \frac{1}{n}$. Therefore the fractional value $\epsilon = \pi/n$ has been found. However, we found satisfactory explanation for the phase difference between “in” and “out” states.

When fermions are coupled to the model, the first problem that appears is the fact that the gauge field has now a short distance behaviour. The fermions eliminate the long range force by means of a screening mechanism. If the fermions are minimally coupled, all fermionic degrees of freedom decouple. On the other hand, the gauge field acquires a mass. This means that a dynamical Higgs mechanism took place. The bosonic particle is no longer confined. The S -matrix of the minimally coupled $\mathbb{C}P^{n-1}$ model confirms that no bound state exists in this case. For the supersymmetric model a similar feature is realized. However, due to the large number of fermionic degrees of freedom, some fermion fields survive the Higgs mechanism. In the case of minimally coupled model, there is no bound state, as one verifies from the absence of poles in the physical region, looking at the solution for the S -matrix describing bosonic scattering, and verified by means of the $1/n$ expansion. The surviving fermionic degrees of freedom are massive, and show a different behavior under chiral transformations as compared with the fermions one started with.

Therefore, in the supersymmetric model, one verifies that the interaction of the gauge field with the fermions reveals a mechanism similar to that involving the interaction of the chiral Gross Neveu model with a gauge field, which assimilates the Goldstone boson of the model, becoming heavy.

In the case of the pure Gross Neveu model, consideration of the physical degrees of freedom leads to a set of $SU(n)$ massive interpolating fields with spin $\frac{1}{2}(1 - \frac{1}{n})$, while the chirality carrying fields decouples. This permits an identification of the antifermion as a bound state of fermions. For the $\mathbb{C}P^{n-1}$ model the combination of the Higgs mechanism for the gauge field with the non-trivial $\mathbb{C}P^{n-1}$ interactions, leads now to a generalization of the mechanism according to which antiparticles are bound states of particles (see (7-9)). As in the interaction of gauge fields with the chiral Gross Neveu fermions, the chirality carrying part of the fermions is absorbed into the gauge field, leaving room for the bound state condition. In the sigma model interaction of $\mathbb{C}P^{n-1}$ model with fermions, the gauge invariant field $\epsilon^{\mu\nu} F_{\mu\nu}$ as well as the chiral density $\bar{\psi}\gamma_5\psi$ get a non-vanishing expectation value in the presence of a θ -term added to the Lagrangian. These first expectation values however vanish in the supersymmetric limit, while the second is maximal, signaling the screening of the external field. This means that the chirality carrying field spurionizes on the gauge invariant subspace, and can not be seen in the physical spectrum.

The first example that some unusual fact is on the way, concerns the example of the $\mathbb{C}P^1$ model. The purely bosonic model is equivalent to the $O(3)$ non-linear σ model. This fact is well established. At the level of S -matrix it is realized by the fact that the $O(3)$ elementary particle is a particle-antiparticle bound state. The bound state pole at the $\mathbb{C}P^1$ level lies at the edge of the physical region, a fact that can be traced back to the confining properties of the model. All these facts have been checked by means of numerical computations¹¹.

In the supersymmetric $\mathbb{C}P^1$ model, the solution of the constraint $\bar{z}_i\psi_i = 0$ can be written as $\bar{\psi}_i = \epsilon_{ij}z_j\chi$, where χ is an anticommuting field. However, from the solution (4-6)

one verifies that χ is empty, and can at most be a Klein factor assuring anticommutativity of the fermion field. This is easily verified at the S -matrix level by means of the identity

$$c_1^{(n=2)}(i\pi - \theta) = \frac{i\pi - \theta}{\theta} \frac{\cos \frac{\theta}{2i}}{\sin \frac{\theta}{2i}} c_1^{(n=2)}(\theta) \quad . \quad (18)$$

With such an identification, it is difficult to reassure supersymmetry! But the worst fact concerns the expected equivalence between $\mathbb{CP}^1$ and $O(3)$ models, which should be realized by the identifications

$$\varphi_i = \sigma_i^{\alpha\beta} z_\alpha \bar{z}_\beta \quad , \quad \text{and} \quad \chi_i = \sigma_i^{\alpha\beta} (z_\alpha \psi_\beta^+ + z_\alpha^+ \psi_\beta) \quad . \quad (19)$$

Absence of fermion-boson backward scattering for the susy $O(3)$ model, is required from the supersymmetric constraints. However it implies that the $d_1(\theta)$ amplitude must vanish.

Let us consider the classical conservation of the supersymmetric current, and latter quantize each step, in order to see, at the end, at which point a different result with respect to the classical case appears. We have

$$J_\mu = \overline{D}_\nu z_i \gamma^\nu \gamma_\mu \psi_i \quad , \quad (20)$$

whose divergence is given by

$$\begin{aligned} \partial^\mu J_\mu &= \overline{D}^2 z_i \psi_i - \frac{1}{2} \epsilon^{\mu\nu} F^{\mu\nu} \bar{z}_i \psi_i + \overline{D} z_i \not{D} \psi_i \\ &= \left(\frac{\alpha}{\sqrt{n}} - m^2 \right) \bar{z}_i \psi_i + \frac{1}{\sqrt{n}} (\bar{\psi}_i c) \psi_i - \overline{D} z_i \left(m + \frac{\varphi}{\sqrt{n}} + i\gamma_5 \varphi_5 \right) \psi_i + \frac{i}{\sqrt{n}} \overline{D} z_i z_i c, \end{aligned} \quad (21)$$

where we used the equations of motion.

The divergence of the supersymmetric current vanishes upon use of the following facts; first one uses a Fierz transformation of the second term. Later one uses the (classical) definitions

$$\sqrt{n} m + \varphi = \frac{f}{\sqrt{n}} \bar{\psi} \psi \quad , \quad \varphi_5 = i \frac{f}{\sqrt{n}} \bar{\psi} \gamma_5 \psi \quad , \quad c = i \frac{2f}{\sqrt{n}} \bar{z} \not{D} \psi \quad , \quad (22)$$

and finally, the constraint $\bar{z}_i \psi_i = 0$. We shall verify that the last equation fails under a very special condition, spoiling conservation of the supersymmetry current.

From this point on we shall use the quantization of the field operators as defined by the BPHZ scheme^{16,17}. Thus, composite field operators are made finite by means of subtractions around zero external momenta.

Identifications of the type (22) are rather subtle in the framework of the $1/n$ expansion, and are related to the fact that for a special vertex $N[\bar{z}\psi]$, the diagram obtained by joining the z and ψ -lines into a $\overline{c}\bar{z}\psi$ vertex, which is the origin of a c -line ending at another vertex of the same type, is cancelled by the diagram where this second vertex originates directly from the $N[\bar{z}\psi]$ vertex. However this is not true when this vertex is oversubtracted.

There are special terms which require special care, arising from the mass term in the equations of motion. Indeed, these terms in principle do not require renormalization, but in order to achieve cancellation of the terms resulting from the equations of motion the renormalization of such terms presents problems.

Indeed, in terms of subtraction of diagrams around zero momenta, one defines various quantum composite operators $N_\delta[\mathcal{O}]$ corresponding to a classical operator $\mathcal{O}$. The index δ defines how a diagram containing the special vertex $N_\delta[\mathcal{O}]$ has to be subtracted^{16,17}. For a diagram with n_ψ external fundamental fermions, n_z external z -bosons, n_c c -lines, etc. we have, for the superficial degree of divergence of a generic diagram (γ) the expression [16,17,18]

$$\delta(\gamma) = 2 - \frac{1}{2}n_\psi - 2n_z - 2n_\alpha - \frac{3}{2}n_c - \cdots + \sum_a (\delta_a - 2) \quad . \quad (23)$$

Moreover the quantum operator $N_\delta[\mathcal{O}]$ satisfies

$$\partial_\mu N_\delta[\mathcal{O}] = N_{\delta+1}[\partial_\mu \mathcal{O}] \quad . \quad (24)$$

A minimally subtracted finite quantum operator has δ equal to the naive dimension of the operator $\mathcal{O}$. This is the least value of δ such that diagrams containing $\mathcal{O}$ be finite. We consider thus the normal product (or quantum operator) $J_\mu = N_{3/2}[\overline{\mathcal{D}}z\gamma_\mu\psi]$. Its divergence is given by terms of dimension 5/2, that is

$$\begin{aligned} \partial^\mu J_\mu &= N_{5/2}[\partial^\mu(\overline{\mathcal{D}}z\gamma_\mu\psi)] \quad , \\ &= N_{5/2}[\overline{D^2}z\psi] + N_{5/2}[\epsilon_{\mu\nu}F^{\mu\nu}\gamma_5\overline{z}\psi] + N_{5/2}[\overline{\mathcal{D}}z\mathcal{D}\psi] \quad . \end{aligned} \quad (25)$$

The last step is to insert the equation of motion inside the normal product. In $1/n$ perturbation this is a standard graphical procedure [17]. Each time one has a factor of the inverse propagator, i.e. $i\mathcal{D} - m$ for fermions, or $\partial^2 + m^2$ for bosons a propagator line is deleted. The resulting diagram is obtained replacing the inverse propagator by the field in the next vertex at which one would arrive following the (deleted) line. In fact, one obtains the terms appearing in the classical equation of motion.

The final step has already been outlined previously and as we mentioned, the step involving the constraint $\overline{z}_i\psi_i = 0$ is dangerous. Indeed, suppose that we have an over-subtraction of this operator, that is, we consider the operator $N_{1/2+\delta'}[\overline{z}_i\psi_i]$ where δ' is a non-negative number.

For the operator we discussed previously we have, from a diagram with one external c -line the contribution

$$N_{5/2}[\overline{z}_i\psi_i]_{1c} = \frac{i}{4\pi m}c - \frac{1}{8\pi m^2}\mathcal{D}c \quad . \quad (26)$$

Since the classical operator $\overline{z}\psi$ is invariant under supersymmetry transformations (2), the above equation signals a non-supersymmetric quantization of this operator.

In fact, such quantization is mandatory. Indeed, to obtain the quantum equations of motion in the way above described, we have to add the mass term, to obtain the inverse propagator, thus we add and subtract to (25) the terms

$$N_{5/2}[m^2\overline{z}\psi] + iN_{5/2}[m\overline{\mathcal{D}}z\psi] \quad . \quad (27)$$

The classical value of the first term, that is $N_{1/2}[m^2\bar{z}\psi]$ is zero, while the second term, $N_{3/2}[m\bar{\mathcal{D}}z\psi]$ is just mc , and together with the term φc will cancel the term $\bar{\psi}\psi$ in the equation of motion of the fermion. Therefore we have to compute the extra subtractions of the terms (27), which are well known in the framework of BPHZ renormalization, corresponding to the so-called Zimmermann identities:

$$N_{\dim(\mathcal{O}+\delta)}[\mathcal{O}] = N_{\dim\mathcal{O}}[\mathcal{O}] + \sum_i r_i N_{\dim(\mathcal{O}+\delta)}[\mathcal{O}_i] \quad , \quad (28)$$

where $\mathcal{O}_i$ are all possible terms with dimension $\dim(\mathcal{O} + \delta)$. The easiest way to compute these terms in the right hand side is to explicitly separate the last (extra) subtractions of the left hand side for each specific diagram. In this way we shall compute r_i , in a $1/n$ perturbativ expansion. We have to compute the extra subtractions.


For the terms at lowest order in $1/n$ we arrive at the partial result

$$\partial^\mu J_\mu = \frac{1}{4\pi} \left[m + \frac{1}{2} \not{\partial} + \frac{1}{\sqrt{n}} (\varphi + i\gamma_5 \varphi_5) \right] c + \dots \quad , \quad (29)$$

where the dots are coming from the diagrams to be considered later.

From the diagram with two external ψ -lines and one external c -line in the figures[†] below. For the first diagram we have:

$$\mathcal{C}_{\alpha\beta,\gamma\delta}^{\Gamma\Gamma'} = m \int \frac{d^2k}{(2\pi)^2} \frac{i}{k^2 - m^2} \left[(k - m) \frac{i}{k - m} \Gamma \right]_{\alpha\beta} \frac{1}{F(k) + i/\pi} \mathcal{O}_{\Gamma\Gamma'} \left(\Gamma' \frac{k + m}{k^2 - m^2} \right)_{\gamma\delta} \quad , \quad (30)$$



Diagrams contributing to the equation (29), with two external fermions. which will contribute with a term of the type $\sum_{\Gamma\Gamma'} \mathcal{C}_{\alpha\beta,\gamma\delta} \psi_\beta \bar{\psi}_\gamma i c_\delta$, where the factor $\mathcal{O}_{\Gamma\Gamma'}$ depends on which of the propagators ($\Delta_\varphi, \Delta_5, \Delta_{\mu\nu}, \Delta_\mu^5$) we are considering. We define the constant

$$\mathcal{W} = m^2 \int \frac{d^2k}{(2\pi)^2} \frac{1}{[F(k) + \frac{i}{\pi}] (k^2 - m^2)^2} \quad . \quad (31)$$

We have the following contributions

- a) $\Gamma = \Gamma' = i$, $\mathcal{O}_{\Gamma\Gamma'} = -1$, $\implies \mathcal{C}_1 = -\mathcal{W} (\bar{\psi} c) \psi$
- b) $\Gamma = \Gamma' = -\gamma_5$, $\mathcal{O}_{\Gamma\Gamma'} = +1$, $\implies \mathcal{C}_2 = -\mathcal{W} \gamma_5 \psi \bar{\psi} \gamma_5 c$
- c) $\Gamma = -\gamma_5, \Gamma' = i\gamma_\mu$, $\mathcal{O}_{\Gamma\Gamma'} = \frac{2mi}{k^2} \epsilon_{\mu\nu} k^\nu$, $\implies \mathcal{C}_3 = +2\mathcal{W} \gamma_5 \psi \bar{\psi} \gamma_5 c$
- d) $\Gamma = i\gamma_\mu, \Gamma' = -\gamma_5$, $\mathcal{O}_{\Gamma\Gamma'} = -\frac{2mi}{k^2} \epsilon_{\mu\nu} k^\nu$, $\implies \mathcal{C}_4 = \mathcal{W} \gamma^\mu \psi \bar{\psi} \gamma_\mu c$

[†] A thin line represents a fermion, a thick one a boson, a dotted-traced line the $c\bar{c}$ propagator, and the wavy line one of the propagators of the remaining auxiliary fields. See [18], or [13] for details.

e) $\Gamma = i\gamma_\mu$, $\Gamma' = i\gamma_\nu$, $\mathcal{O}_{\Gamma\Gamma'} = -\left(g_{\mu\nu} - \frac{k_\mu k_\nu}{k^2}\right)$, $\implies \mathcal{C}_5 = -\frac{1}{2}\mathcal{W}\gamma^\mu\psi\bar{\psi}\gamma_\mu c$.
Adding all these contributions and performing a Fierz transformation, we have

$$\sum \mathcal{C} = \mathcal{W} \left[\frac{1}{2}\bar{\psi}\psi - \frac{1}{2}\bar{\psi}\gamma_5\psi\gamma_5 + \bar{\psi}\gamma^\mu\psi\gamma_\mu \right] c \quad . \quad (32)$$

Consider now the second diagram. It is given by

$$\mathcal{E}_{\alpha\beta,\gamma\delta} = i^2 \int \frac{d^2k}{(2\pi)^2} \left(\frac{2(\not{k} - 2m)}{F(k) + i/\pi} \right)_{\alpha\beta} \left(\frac{i}{k^2 - m^2} \right)^2 \left(\frac{i}{\not{k} - m} \right)_{\gamma\delta} , \quad (33)$$

which will contribute as

$$\mathcal{E} = \mathcal{E}_{\alpha\beta,\gamma\delta} \bar{\psi}_\alpha \psi_\beta i c_\delta \quad . \quad (34)$$

Defining now

$$\mathcal{W}' = m^2 \int \frac{d^2k}{(2\pi)^2} \frac{1}{[F(k) + i/\pi](k^2 - m^2)^3} \quad (35)$$

we find

$$\mathcal{E} = \mathcal{W}' [4\bar{\psi}\psi - \bar{\psi}\gamma^\mu\psi\gamma_\mu] c \quad . \quad (36)$$

Therefore we have

$$\left[\left(\frac{1}{2}\mathcal{W} + 4\mathcal{W}' \right) \bar{\psi}\psi + (\mathcal{W} + \mathcal{W}') \bar{\psi}\gamma_\mu\psi\gamma^\mu - \frac{1}{2}\mathcal{W}\bar{\psi}\gamma_5\psi\gamma_5 \right] c \quad . \quad (37)$$

This equation can be used to choose the coupling constant in $\sqrt{n}m + \varphi = \frac{f}{\sqrt{n}}\bar{\psi}\psi$ in order to cancel the φ contribution to the right hand side of equation (29), but not the φ_5 and $\bar{\psi}\gamma_\mu\psi$ contributions at the same time. Therefore we arrive at

$$\partial^\mu J_\mu = \mathcal{W}_1 \bar{\psi}\gamma_5\psi\gamma_5 c + \mathcal{W}_2 \bar{\psi}\gamma_\mu\psi\gamma^\mu c \quad , \quad (38)$$

which we claim to be a candidate to the supersymmetry anomaly. The detailed computation of $\mathcal{W}_1$ and $\mathcal{W}_2$ depends on the normalization of the current.

In fact, one expects that one of the supersymmetry currents be conserved, since the model can be formulated in an explicitly supersymmetric fashion¹⁹.

A further indication that the supersymmetric $\mathbb{C}P^{n-1}$ model has a higher symmetry preventing the boson fermion back scattering amplitude to be vanishing when the constraint $\bar{z}_i\psi_i = 0$ is strongly realized, comes from the perturbative computation. Indeed, up to fourth order in a perturbative computation, there is no contribution for $d_1(\theta)$ in (3).

We have verified that the S -matrix of the $\mathbb{C}P^{n-1}$ model is supersymmetric if one considers non trivial phase factors for asymptotic states. This may be a consequence of a possible supersymmetry anomaly which would appears due to the quantization of the constraint $\bar{z}_i\psi_i = 0$. Nevertheless, this can also be understood from further properties of the model. Indeed, the implicit Higgs mechanism shifting the pole of a gauge field away from the origin, implies a set of bound states connecting tightly particles and antiparticles,

and more importantly, fermions and bosons. Therefore the statistics got lost. The Klein factor used in two-dimensions to reassure the correct statistics after bosonization does not seem to be enough for a rebuilding of the supersymmetry algebra.

Although the bound state structure might be seen at first look as a consequence of supersymmetry, generalizing the structure of the chiral Gross Neveu model, a concrete realization of supersymmetry at the level of the S -matrix fails if statistics is not taken into account. The supersymmetry algebra, if reconstructed, should act on the fields with non trivial phase factors due to asymptotic statistics.

Consequences for higher dimensional physics are vast. In three dimensions¹⁹ there are two phases in the model, and in one of them the issue of supersymmetry displays a behavior similar to the one presented here²⁰. This opens the possibility of describing particles at very high energy in terms of supersymmetric grand unified theories, while at sufficiently low energies supersymmetry breaks as a consequence of dynamics.

The fact that composite operators might be responsible for supersymmetry breaking has been conjectured before²¹. In fact some candidates have been written for such composite (anomalous) operators. Although we did not check in detail the relation to this line of works, it is conceivable that both issues can be related.

Several consistency checks of the results have been performed and will be reported elsewhere. First the result is non-perturbative. We have checked that d_1 is zero up to fourth order in the coupling constant, while being non zero in the $1/n$ expansion. Moreover the break of supersymmetry is minimal, in the sense that at lowest order in $1/n$ the commutator of the supersymmetry charge with the S operators is a constant, which is presumably necessary in order to avoid further problems. In the $\mathbb{CP}^1$ case the d term in the S -matrix may imply non equivalence with the supersymmetric $O(3)$ model. Moreover, the algebra underlining the bound state structure does not seem to accomodate supersymmetry transformations in a simple way.

Finally, we should also point out that consistency of the supersymmetry transformations, $U(1)$ current conservation and soliton type boundary condition for the auxiliary fields as initially pointed out by Witten¹³, is consistent with our results. A supersymmetry transformation of the naive $U(1)$ current, $J_\mu = i\bar{z}\partial_\mu z + \bar{\psi}\gamma_\mu\psi$ leads to an expression of the type

$$-\frac{1}{2\sqrt{n}}\bar{c}\gamma_\mu\epsilon - \frac{1}{2\sqrt{n}}\bar{\epsilon}\gamma_\mu c - \frac{i}{2}\bar{J}_\mu^{susy}\epsilon + \frac{i}{2}\bar{\epsilon}J_\mu^{susy}$$

We expected this expression to be conserved. On the other hand, c is in the same supermultiplet as the remaining auxiliary fields, whose boundaries change by phases¹³ $e^{\frac{i\pi}{n}}$. The above association with the supersymmetry current in order to achieve conservation would imply such phases in the asymptotic values of the fields and charges. In fact, the low energy effective equation of motion for c implies a term of the form $\frac{i}{4\pi}\phi_5\gamma_5 c$, leading to contributions of the type $\bar{\psi}\gamma_5\psi\gamma_5$ and $\bar{\psi}\gamma_\mu\psi\gamma^\mu$ after use of the Ward identity. For such boundary conditions, the contribution from a supersymmetry current to the Wess Zumino consistency condition, seems to be unavoidable. These issues are currently under investigation.

Acknowledgements: the authors would like to thank L. Alvarez-Gaumé, D. Freedman, O. Piguet, K. Sibold, and especially R. Stora for discussions.

References

- [1] D. Gepner, Commun. Math. Phys. **141** (1991)381; Nucl. Phys. **B296** (1987)380
- [2] C. Vafa, Mod. Phys. Lett. **A4** (1989)1169
- [3] P. Fendley, S.D. Mathur, C. Vafa and N. Warner, Phys. Lett. **283B** (1990)257
- [4] C. Vafa and N. Warner, Phys. Lett. **B218** (1989)51
E. Martinec, Phys. Lett. **217B** (1989)431
- [5] V.I. Arnold, S.M. Guseian-Zadeh and A.N. Varchenko, *Singularities of Differentiable Maps*, Birkhauser, 1985
- [6] P. Fendley and K. Intriligator, Nucl. Phys. **B372** (1992)533
- [7] W. Lerche, C. Vafa and N. Warner, Nucl. Phys. **B324** (1989)427
- [8] Y. Kazama, H. Suzuki, Mod. Phys. Lett. **A4** (1989)235; Phys. Lett. **B216**(1989)112; Nucl. Phys. **B321**(1989)232.
- [9] R. Köberle and V. Kurak, Phys. Lett. **191B** (1987)295
- [10] M.C.B. Abdalla and A. Lima Santos, Acta Phys. Pol. **B15** (1984)813
M. Karowski, V. Kurak and B. Schroer, Phys. Lett. **81B** (1979)200
- [11] E. Abdalla, M.C.B. Abdalla, N. Alves, C.E.I. Carneiro, Phys. Rev. **D41** (1990)571
- [12] O. Piguet, and K. Sibold, Nucl. Phys. **B253**(1985) 269; P.L. White, Class. Q. Gravity **9**(1992) 413,1663
- [13] A. D’Adda, P. diVecchia and M. Lüscher, Nucl. Phys. **B152** (1979)125
E. Witten, Nucl. Phys. **B149** (1979)285
- [14] R. Köberle and V. Kurak, Phys. Rev. **D36** (1987)627
- [15] E. Witten and D. Olive, Phys. Lett. **78B** (1978)97
- [16] N.N. Bogoliubov and O.S. Parasiuk, Acta Math. **97** (1957)227
K. Hepp, Commun. Math. Phys. **2** (1966)301
W. Zimmermann, *Lectures on Elementary Particles and Quantum Field Theory*, Vol 1, 1970, Brandeis Univ. Summer Institute.
- [17] J. Lowenstein, BPHZ Renormalization; Lectures given at Int. School of Mathematical Physics, Erice, Sicily, August 17-31, 1975, Erice Math. Phys. 1975:95, (QCD161:I651:1975).
- [18] E. Abdalla, M.C.B. Abdalla and K. Rothe, *Non Perturbative Methods in 2D Quantum Field Theory*, World Scientific Publishing, 1991
- [19] P. Gaigg, M. Schweda, O. Piguet and K. Sibold, Fortschritte der Physik **32**(1984) 623
- [20] E. Abdalla and M.C.B. Abdalla, work in progress
- [21] J.A. Dixon, Talk given at the Harc conference on recent advances in the superworld, hep-th/9308088